



YaraVita™

Zintrac® 700



5 litres
Net weight
8.67 kg

NOTICE

Before using this product, read the entire label

Zintrac® 700

UK FERTILISER

Zinc oxide fertiliser suspension

Guaranteed analysis % w/w:

40.9% (=709 g/l) Zinc (Zn) as oxide, total

To be used only where there is a recognised need.

Do not exceed the application rate

INFORMATION FOR ROI MARKET

PFC 1(C)(II)(a) Straight Inorganic micronutrient fertilizer
Liquid Suspension

Content % w/w:

40.9% (=709 g/l) Zinc (Zn) as oxide, total

Also Contains % w/w:

1% Total Nitrogen (N)

[1% Urea Nitrogen]

Ingredients: Zinc oxide CAS n° 1314-13-2 (CMC 1: Virgin material substances and mixtures), monoethylene glycol CAS n° 107-21-1 (CMC 1).

To be used only where there is a recognised need. Do not exceed the application rate. This fertiliser contains urea, which can release ammonia and have an impact on air quality. Depending on local conditions, appropriate remediation measures must be taken.

Batch Number: See Pack

Manufacture Date: See Pack

Produced and Distributed by: **Yara UK Limited.**

Pocklington Industrial Estate, Pocklington, York. YO42 1DN

Tel: 01759 302545 www.yara.co.uk

ROI: Imported by: **Yara UK Ltd.** The Old Rectory, Cork Road, Carrigaline, P43 W921. County Cork.

Tel: +44 1472 889 250 www.yara.ie

YaraVita™ Zintrac®700

APPLICATION RATES AND TIMINGS

Apple, Pears: 1 to 2 l/ha at bud burst and at the end of flowering. Avoid treatment during flowering. A further application of 1 to 2 l/ha should be made after harvest but before leaf senescence. Water rate: 200-1,000 l/ha.

Carrot: 1 l/ha when crop is 15 cm tall. For moderate to severe deficiency repeat applications at 10 to 14 day intervals. Water rate: 200 l/ha.

Cereals: 1 l/ha from 2 leaf stage to second node detectable (Zadok's G.S. 12 to 32). Also, 0.5 l/ha from ear fully emerged to the end of flowering (Zadok's G.S. 59 to 69). Water rate: 50 to 200 l/ha. Seed Dressing: Typical rate: 3 litres per tonne. Increase up to 8 litres per tonne if necessary.

Cherry, Plum: 1 to 2 l/ha applied at winter bud and again at pink bud and again after harvest but before leaf senescence. Water rate: 500 to 1,000 l/ha.

Conifers: 2 applications of 1 l/ha once there is new season leaf production and again in early autumn. Water rate: 500 to 1,000 l/ha.

Grass (Amenity): 1 l/ha as soon as growth commences in spring and/or following identification of need by analysis. Repeat sprays at 10 to 14 day intervals as necessary. Water rate: 200 l/ha.

Grass (Grazing): 1 l/ha 10 to 14 days prior to turnout. Animals should be kept out of treated pastures for 10 days to allow for nutrient uptake by the herbage. Water rate: 200 l/ha

Grass (Silage/Hay): 1 l/ha when crop is 15 cm tall. For severe deficiency repeat applications should be made at the above rate at the 15 cm stage before each subsequent cut. Water rate: 200 l/ha.

Hops: 1.5 l/ha when crop is 2.5 metres tall. Water rate: 500 l/ha.

Leek: 1 l/ha applied two weeks after transplanting, or in the case of direct sown crops, when the crop is 15 cm tall. One or two repeat applications may be necessary at 10 to 14 day intervals. Water rate: 200 l/ha.

Linseed: 1 l/ha when crop is 2 to 15 cm tall. Water rate: 200 l/ha.

Maize: 1 l/ha from 3 to 8 leaf stage. For severe deficiency, repeat application should be made at 10 to 14 day intervals. Water rate: 200 l/ha.

Onion: 1 l/ha when there is sufficient leaf area to intercept the spray. Water rate: 200 l/ha.

Peas: 0.5 to 1 l/ha when crop is 5 to 15 cm tall. Water rate: 50 to 200 l/ha.

Potatoes: 1 l/ha one week after 100% emergence. For moderate to severe deficiency, repeat applications may be necessary at 10 to 14 day intervals. Also, 1 l/ha following petiole analysis, during tuber bulking. Water rate: 200 l/ha.

Rasperry: 0.5 at green bud. Water rate: 200 to 500 l/ha.

Strawberry (Field Grown): One application of 0.5 l/ha at green bud followed by two applications of 0.25 l/ha applied at white bud and post-harvest regrowth. Water rate: 200 to 500 l/ha.

Sugar Beet: 1 l/ha at 4 to 6 leaf stage. For moderate to severe deficiency, repeat at 10 to 14 day intervals. Water rate: 200 l/ha.

Turf: 1 l/ha as soon as growth commences in spring and/or following identification of need by analysis. Repeat sprays at 10 to 14 day intervals as necessary. Water rate: 200 l/ha.

PROTECTED CROPS

0.05 litres per 100 litres water maximum concentration. Water rate: 1000 l/ha maximum. Refer to equivalent field grown crop for application timing.

For further advice and product information please scan QR code using your smart phone.



UK



ROI

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use. Please refer to the MSDS for further information, available at www.yara.com.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from temperature extremes (-5°C < and <35°C). Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



YaraVita™ Zintrac® 700



Warning

H410: Very toxic to aquatic life with long lasting effects.

P273: Avoid release to the environment.

P391: Collect spillage.