

# YaraVita Bud Builder FL

## APPLICATION RATES AND TIMINGS

**Apple, Pears:** 10 l/ha applied post-harvest before leaf fall. Water rate: 500 to 1000 l/ha.

**Blackcurrant, Boysenberries, Raspberry, Strawberry (field grown):** 5 l/ha applied from green bud to just prior to start of flowering. Also, 5 l/ha post-harvest, pre-leaf senescence. Water rate: 200 to 500 l/ha.

**Apricot, Cherry, Nectarines, Peach, Plum:** 10 l/ha applied post-harvest before leaf fall. Repeat if necessary leaving 10 to 14 days between applications. Water rate: 500 to 1000 l/ha.

**Citrus:** 10 l/ha applied post-harvest. Also, 5 to 10 l/ha applied from bud burst to start of flowering. N.B. Do not apply during flowering. Water rate: 500 l/ha.

**Cucurbits (field grown):** 5 l/ha at the 4 to 6 leaf stage. Repeat applications may be necessary at 10 to 14 day intervals. Water rate: 500 l/ha. N.B. Do not apply during flowering.

**Grapevines:** 1 to 3 applications of 5 l/ha applied pre-flowering at a 10-14 day interval. Also, 10 l/ha applied post-harvest before leaf fall. Water rate: 500 to 1000 l/ha  
**Kiwifruit:** 5 to 10 l/ha applied pre-flowering. Also, 5 to 10 l/ha applied post-harvest before leaf fall. Water rate: 500 to 1000 l/ha.

**Nuts (deciduous):** 10 l/ha applied post-harvest before leaf fall. Also, 5 to 10 l/ha applied from bud burst to start of flowering. Water rate: 500 to 1000 l/ha.

**Olive:** 5 to 10 l/ha applied from bud burst to start of flowering. Water rate: 500 l/ha.

**Potatoes:** 3 l/ha one week after 100 percent emergence. Repeat once or twice at 10 to 14 day intervals for moderate to severe deficiency. Water rate: 200 L/ha.

**Tomato:** 1 to 3 l/ha at the 4 to 6 leaf stage. One or two repeat applications may be necessary at 10 to 14 day intervals. N.B. Do not apply during flowering. Water rate: 500 l/ha.

## DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS. Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

## TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit [www.tankmix.com](http://www.tankmix.com) for important information. Alternatively, contact Yara for specific advice.

## PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

## STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes (temperatures exceeding 40°C). Dispose of contents and container in accordance with all local, regional, national and international regulations.

## CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

