



YaraVita™ Caltrac™



1 liter

NOTICE

Before using this product, read the entire label

Safety data sheet available for professional user on request

SHAKE WELL BEFORE USE

EC FERTILIZER

Calcium-based fertilizer

Guaranteed analysis:

16% w/v = 160 g/l Calcium (Ca) total

FPA Registration No. 1-IIF-3499

Manufactured by: Yara UK Ltd, Pocklington, York, YO42 1DN
Distributed by: **Yara Fertilizers Philippines Inc.**
1605 One Global Place, 5th Avenue cor. 25th St.,
Bonifacio Global City, Taguig City, Philippines
Tel. Nos.: (02) 8845-3801 to 05

YaraVita Caltrac™

Crop	Remarks
Eggplant	Spray young fruits directly with a solution containing 0.4 % Caltrac (400 ml in 100 l water). Alternatively: Two to four applications of 5 l/ha from the end of flowering at 7 day intervals. Water rate: 200 to 500 l/ha.
Pepper (Field Grown)	Spray very young fruits directly, using 400 ml per 100 liters of water or 66 ml per 16 liters water. Maximum water rate: 1,000 l/ha. Alternatively: Regular applications of 5 l/ha from flowering on second truss onwards. Water rate: 50 to 500 l/ha For knapsack sprayer application: 160 ml per 16 liters water (Do not exceed 5 l/ha product per application).
Potatoes	2 to 3 applications of 5 to 10 l/ha starting at tuber initiation (50% of the tip swelling are twice the diameter of the rest of the stolon) and following petiole analysis during tuber bulking with 10 to 14 day application interval. Water rate: 200 l/ha. For knapsack sprayer application: 400 ml to 800 ml per 16 liters water (Do not exceed 10 l/ha product per application).
Tobacco	5 l/ha applied two to three weeks after transplanting (3 to 4 leaf stage) and again 10 days later. Water rate: 30 to 500 l/ha. For knapsack sprayer application: 400 ml per 16 liters water (Do not exceed 5 l/ha product per application).
Tomato (Field Grown)	Application of 5 l/ha commencing from flowering on second truss. Repeat application at 7 to 14 days interval. Water rate: 50-500 l/ha. Alternatively: Spray directly on very young fruits using 400 ml per 100 liters water. Maximum water rate: 1,000 l/ha. For knapsack sprayer application: 160 ml per 16 liters water (Do not exceed 5 l/ha product per application).

10ml = 1 tbsps

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read all labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 40°C) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

Warning

Hazardous ingredients: calcium chloride

H319: Causes serious eye irritation.

P280: Wear eye protection.

P264: Wash hands thoroughly after handling.

P305+P351+P338+P337+P313: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical attention.

National Chemical Emergency Council Call: **02-8231-21-49**

