

YaraVita Thiotrac™ 300

APPLICATION RATES AND TIMINGS

Citrus: Two applications of 1 quart/acre at a 10 to 14 day interval before flowering. Water rate: 50 gallons/acre.

Cotton: 0.5 to 1 quart/acre one month after 100% emergence. Repeat as required at 10 to 14 day intervals.

Avoid flowering. Water rate: 20 gallons/acre

Maize: 1 quart/acre at the 4 to 8 leaf stage. Repeat 10 to 14 days later if necessary.

Water rate: 3 to 20 gallons/acre.

Potatoes: 1 quart/acre one week after 100 percent emergence. Repeat applications may be necessary at 10 to 14 day intervals. Also, 1 quart/acre following petiole analysis during tuber bulking, and at 10 to 14 day intervals during tuber bulking if necessary. Water rate: 5 to 20 gallons/acre.

Soya bean: 1 to 2 quarts/acre at the 4 to 6 leaf stage. Repeat at 10 day intervals if required.

Water rate: 20 gallons/acre.

Wheat: 1 to 2 quarts/acre at pseudo stem erection to first node detectable (Zadok's G.S. 30 to 31). For moderate to severe deficiency, repeat applications at 10 to 14 day intervals. Water rate: 3 to 20 gallons/acre.

Alternatively: 1 to 2 applications at 2 to 3 pints/ac from tillering to 1st node detectable at a 10 to 14 day interval.

Water rate: 20 gallons/acre. Wheat only: Up to 2 quarts/acre from ear fully emerged to the end of the milk development stage in grain ripening (Zadok's G.S. 59 to 79) or two applications of 1 quart/acre in the period flag leaf ligule just visible to the end of the milk development stage in grain ripening (Zadok's G.S. 39 to 79). Water rate: 3 to 20 gallons/acre.

Extended recommendations available at <https://www.yara.us/crop-nutrition/fertilizer-products/yaravita/>

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 100°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



WARNING: Cancer and Reproductive Harm –www.P65Warnings.ca.gov