



YaraVita™

Molytrac™
250



1 litre

NOTICE

Before using this product, read the entire label

GROUP 2 FERTILIZER
REGISTRATION No. B3709 Act 36 of 1947

Molybdenum fertilizer in solution for
application to deficient crops

Guaranteed analysis - soluble in water:
155 g/kg = 250 g/l molybdenum (Mo)

Typical density 1.61 kg/litre at 20°C

Manufactured by Yara UK Ltd., Pocklington, York
YO42 1DN, U.K.
Registered by Yara Africa Fertilizers (PTY) Ltd.,
Reg.nr. 2011/003434/07
The Pivot @ Montecasino, No. 1 Montecasino
Boulevard, Fourways, Johannesburg 2191.

RECOMMENDED APPLICATION RATES AND TIMINGS

Brassicas: 250 to 300 ml/ha from 4 to 6 leaf stage. For moderate to severe deficiency, repeat applications should be made at the above rate at 10 to 14 day intervals. Water rate: 50-500 l/ha.

Citrus: 0.25 l/ha from 2/3 of new leaf development in the spring. For moderate to severe deficiency repeat application 10 to 14 days later. Also, 0.25 l/ha applied during autumn flush. Repeat 10 to 14 days later if necessary. Water rate: 500 l/ha.

Cucurbits (Field Grown): 250 ml/ha 1 to 2 weeks after germination and at 4 leaf stage. Water rate: 50-200 l/ha.

Grapes: 0.5 l/ha applied before flowering. Repeat at fruit set, bunch closure and early veraison if necessary. Water rate: 200 l/ha.

Legumes, Pulses: 250 to 500 ml/ha when plants are 5 to 15 cm tall. For moderate to severe deficiency, repeat applications should be made at the above rate at 10 to 14 day intervals. Water rate: 50 to 200 l/ha.

Lucerne: 250 to 500 ml/ha every 2 to 3 cuts. Water rate: 50-200 l/ha.

Maize: 0.25 to 0.5 l/ha at 6-8 leaves. Water rate: 50-200 l/ha.

Peas: 250 to 500 ml/ha when plants are 5 to 15 cm tall. For moderate to severe deficiency, repeat applications should be made at the above rate at 10 to 14 day intervals. Water rate: 50 to 200 l/ha.

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes (temperatures exceeding 40°C). Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



26.03.21

WILZAYP87M00