

# YaraVita B-Moly™

## APPLICATION RATES AND TIMINGS

### WARNING

This product contains Boron (B) and Molybdenum (Mo) which may be injurious to certain crops and toxic to ruminant animals. Contact your county agent or field consultant for specific information.

**Cotton:** 1-2 pints/acre at 4 to 6 leaf stage, at appearance of first flower bud squares and again at open flowers stage.

Water rate: 5 to 15 gallons/acre.

**Maize:** Corn: 1-2 pints/acre at 4 to 8 leaf stage. For moderate to severe deficiency, a repeat application may be necessary 10 to 14 days later. Water rate: 10 to 20 gallons/acre.

**Potatoes:** One to two applications of 1-2 pints/acre applied from 7 to 14 days after 100 percent emergence to 20 days after tuber initiation and, following petiole analysis, during tuber bulking. Water rate: 10 to 20 gallons per acre.

**Soybeans:** 1-2 pints/acre when crop is 2 to 6 inches tall, repeated at 10 to 14 day intervals if necessary.

Water rate: 5 to 20 gallons/acre.

*Extended recommendations available at <https://www.yara.us/crop-nutrition/fertilizer-products/yaravita/>*

---

### DIRECTIONS FOR USE

#### ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

### TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit [www.tankmix.com](http://www.tankmix.com) for important information. Alternatively, contact Yara for specific advice.

### PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

### STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 100°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

### CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.



**WARNING:** Cancer and Reproductive Harm –[www.P65Warnings.ca.gov](http://www.P65Warnings.ca.gov)